

MEETING ABSTRACT

Open Access

Endoscopic intranasal orbital decompression in the course of Graves' orbitopathy

Antoni Krzeski*, Mirosław J Szczepański

From 4th Congress of the Polish Thyroid Association 2013
Lodz, Poland. 11-13 April 2013

The authors present the endoscopic intranasal orbital decompression, one of the surgical techniques used for orbital decompression. It can be applied for various indications. The most common indications are: maxillo-facial trauma, orbital haemorrhage, subperiorbital abscess as the complication of sinusitis, optic neuropathy, thyroid-associated orbitopathy (Graves' orbitopathy). The authors conclude that the endoscopic transnasal orbital decompression is a valuable and safe operational technique for patients requiring orbital decompression.

Published: 5 April 2013

doi:10.1186/1756-6614-6-S2-A66

Cite this article as: Krzeski and Szczepański: Endoscopic intranasal orbital decompression in the course of Graves' orbitopathy. *Thyroid Research* 2013 **6**(Suppl 2):A66.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Department of Otolaryngology, Medical University of Warsaw, Warsaw, Poland



© 2013 Krzeski and Szczepański; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.